

Singapore Management University

Institutional Knowledge at Singapore Management University

Research Collection School Of Computing and
Information Systems

School of Computing and Information Systems

5-1997

A measurement of Rb using a lifetime-mass tag

R. BARATE

Manoj THULASIDAS

Singapore Management University, manojt@smu.edu.sg

Follow this and additional works at: https://ink.library.smu.edu.sg/sis_research



Part of the [Physics Commons](#)

Citation

BARATE, R. and THULASIDAS, Manoj. A measurement of Rb using a lifetime-mass tag. (1997). *Physics Letters B*. 401, (1-2), 150-162.

Available at: https://ink.library.smu.edu.sg/sis_research/3668

This Journal Article is brought to you for free and open access by the School of Computing and Information Systems at Institutional Knowledge at Singapore Management University. It has been accepted for inclusion in Research Collection School Of Computing and Information Systems by an authorized administrator of Institutional Knowledge at Singapore Management University. For more information, please email cherylds@smu.edu.sg.

A Measurement of R_b using a Vertex Mass Tag

The SLD Collaboration*

Stanford Linear Accelerator Center

Stanford University, Stanford, CA 94309

Submitted to Physical Review Letters

Abstract

We report a new measurement of $R_b = \Gamma_{Z^0 \rightarrow b\bar{b}}/\Gamma_{Z^0 \rightarrow \text{hadrons}}$ using a double tag technique, where the b hemisphere selection is based on the reconstructed mass of the B hadron decay vertex. The measurement was performed using a sample of 130k hadronic Z^0 events, collected with the SLD at the SLC. The method utilizes the 3-D vertexing abilities of the CCD pixel vertex detector and the small stable SLC beams to obtain a high b -tagging efficiency and purity. We obtain $R_b = 0.2142 \pm 0.0034(\text{stat.}) \pm 0.0015(\text{syst.}) \pm 0.0002(R_c)$.

We report a new measurement of R_b , the fraction of $Z^0 \rightarrow b\bar{b}$ events in hadronic Z^0 decays, collected with the SLD at SLC using a mass tag technique. The ratio R_b is of special interest as a test of the Standard Model (SM), since it is sensitive to possible new physics effects which modify the radiative corrections to $Zb\bar{b}$ vertex. The vertex corrections are isolated because R_b is a ratio between two hadronic rates, hence propagator (oblique), radiative and QCD corrections common to all quark flavors mostly cancel. Recent measurements yielded a world average R_b value 3σ higher than that predicted by the SM [1]. Previous measurements [2] selected $b\bar{b}$ events based upon mainly the long B hadron lifetime and were limited systematically by contamination in the sample from residual $c\bar{c}$ events. To avoid this limitation our b -tag exploits the large B mass, since the mass distribution has a very small charm contamination beyond the charm mass cutoff. Taking advantage of SLD's precise 3-D vertexing capability and the small and stable SLC beam spot, we achieve a very efficient and pure b selection. We use a self-calibrating double tag technique [2], which allows one to measure both R_b and the b -tag efficiency, ϵ_b , simultaneously.

This measurement is performed using approximately 130K $e^+e^- \rightarrow Z^0 \rightarrow q\bar{q}$ events collected during 1993-95. A detailed description of the detector can be found elsewhere [3]. We used the information from charged particle tracks measured with the CCD pixel Vertex Detector (VXD) along with the Central Drift Chamber. The event selection and the determination of the thrust axis use the the energy deposits measured with the Liquid Argon Calorimeter.

The luminous region of the SLC interaction point (IP) has a size of about $(1.5 \times 0.8) \mu\text{m}$ in the x,y plane transverse to the beam direction and $700 \mu\text{m}$ along the beam direction. We use the average IP position of small groups of sequential hadronic events to determine the primary vertex (PV) in the x-y plane. The longitudinal position of the PV is determined for each event individually [3]. This results in a PV position measurement with uncertainties of $7 \mu\text{m}$ transverse to the beam axis and $35 \mu\text{m}$ ($52 \mu\text{m}$ for $b\bar{b}$ events) along the axis. The measured track impact parameter resolution is $\sigma_{r\phi}[\mu\text{m}] = 11 \oplus 70/p \sin^{3/2} \theta$, $\sigma_{rz}[\mu\text{m}] = 37 \oplus 70/p \sin^{3/2} \theta$ where p is the track momentum expressed in GeV/c .

The hadronic event selection is based on charged track multiplicity and track visible energy requirements as described in Ref. [3]. The event selection is studied with Monte Carlo (MC) events generated using JETSET 7.4 event generator [4], where the B hadron decays are simulated using a model tuned to current B and D decay data [5]. A plane transverse to the thrust axis is used to divide the event into two hemispheres. In order to ensure that the events are well contained within the acceptance of the VXD, the polar angle of the thrust is required to be within $|\cos \theta_{thrust}| < 0.71$. In addition, to ensure the event hemisphere division is sensible and to reduce the contribution from events containing $g \rightarrow b\bar{b}$, we require that the event contain no more than three jets (defined using charged tracks and the JADE algorithm [6] with $y_{cut}=0.02$). A total of 72074 events were selected.

In each event well measured tracks [3] are used to search for a secondary vertex (SV). The SV are found by searching for areas of high track overlap density from the individual track resolution functions, in 3-D co-ordinate space [7]. The SV are required to be separated from

the PV by at least 1 mm and to contain at least two tracks each with a 3-D impact parameter with respect to the $IP \geq 130\mu\text{m}$, ensuring they originate from the decay of a particle with relatively long lifetime. Simulation studies show that secondary vertices are found in 50% of all b hemispheres, in 15% of the charm and $< 1\%$ of the light quark hemispheres [7].

Due to the cascade structure of the B decay, not all the tracks in the decay chain will come from a common decay point, thus the SV is incomplete. We improve our estimate of the B decay vertex mass by attaching to the SV additional tracks that are consistent with the hypothesis of originating from the same SV . We illustrate this in Fig. 1(a). We define the vertex axis to be the straight line between the PV and SV centroids. For each track not in the SV the 3-D distance of closest approach, T , and the distance from the PV along the vertex axis to this point, L , are calculated. Tracks with $T < 1$ mm and $L/D > 0.25$, where D is the distance from the PV to the SV are attached to the SV to form a B decay candidate. The invariant mass, M_{ch} , of the B candidate is obtained assuming each track has the mass of a charged π ; the distribution of M_{ch} is shown in Fig. 2(a). We require M_{ch} to be well above the charm mass, $M_{ch} > 2$ GeV/ c^2 , results in a b hemisphere tagging efficiency of 28% with a purity of 98%.

We improve the b tagging efficiency by applying a kinematic correction to the calculated M_{ch} . Due to the neglect of information about the neutral particles in the decay, the SV flight path and the SV momentum vector are typically acollinear. In order to compensate for the acollinearity we correct M_{ch} using the minimum missing momentum (P_t) transverse to the SV flight path. To reject non- $b\bar{b}$ events with an artificially large P_t due to detector resolution effects, we define P_t with respect to a vector tangent to the error boundaries of both the PV and the SV , such that P_t is minimized (see Fig. 1(b)). The ability to make this minimal correction is most effective at SLD due to the small and stable beamspot of the SLC and the high resolution vertexing. We then define the P_t -corrected mass, $\mathcal{M} = \sqrt{M_{ch}^2 + P_t^2} + |P_t|$, and require $\mathcal{M} \leq 2 \times M_{ch}$ to reduce the contamination from fake vertices in light quark events. The distribution of \mathcal{M} is shown in Fig. 2(b). By requiring $\mathcal{M} > 2\text{GeV}/c^2$ we significantly raise our b -tag efficiency, yielding $\epsilon_b = 35.3\%$ for the same purity.

We measure R_b and ϵ_b by counting the fraction of the event sample containing one tagged hemispheres, F_s , and the fraction containing both hemispheres tagged, F_d :

$$R_b = \frac{(F_s - R_c(\epsilon_c - \epsilon_{uds}) - \epsilon_{uds})^2}{F_d - R_c(\epsilon_c - \epsilon_{uds})^2 + \epsilon_{uds}^2 - 2F_s\epsilon_{uds} - \lambda_b R_b(\epsilon_b - \epsilon_b^2)},$$

$$\epsilon_b = \frac{F_d - R_c\epsilon_c(\epsilon_c - \epsilon_{uds}) - F_s\epsilon_{uds} - \lambda_b R_b(\epsilon_b - \epsilon_b^2)}{F_s - R_c(\epsilon_c - \epsilon_{uds}) - \epsilon_{uds}}.$$

The only term dependent upon B production and decay modeling is the b hemisphere tagging correlation, $\lambda_b = \frac{\epsilon_b^{double} - \epsilon_b^2}{\epsilon_b - \epsilon_b^2} = 0.59\%$, where we have used the simulation to estimate λ_b . Estimates of the hemisphere tagging rates of light quarks, $\epsilon_{uds} = 0.06\%$ and charm quarks, $\epsilon_c = 0.69\%$, are also derived from the simulation, and we assume $R_c = \frac{\Gamma_{Z^0 \rightarrow c\bar{c}}}{\Gamma_{Z^0 \rightarrow q\bar{q}}} = 0.171$. We measure $R_b = 0.2142 \pm 0.0034_{stat}$, which includes a correction of $+0.0003$ for the $e^+e^- \rightarrow \gamma \rightarrow b\bar{b}$ contribution as calculated by ZFITTER [8]. The measured value of $\epsilon_b = 35.3\% \pm 0.6\%$ is in good agreement with the MC estimate of 35.5%. As a cross-check we repeated the

measurement using different \mathcal{M} cuts; the results are summarized in Fig. 3(a). The R_b results are consistent for values of $0 < \mathcal{M}_{cut} < 3 \text{ GeV}/c^2$.

The systematic uncertainty on R_b , given in detail in Table 1, results from a combination of detector related effects and physics uncertainties in the simulation which affect our estimates of ϵ_c , ϵ_{uds} and λ_b . The physics systematic errors are assigned by comparing the nominal simulation distributions with an alternative set of distributions which reflect the uncertainties in the world average measurements of the MC physics parameters [9]. The two significant sources of systematic errors from light quark events come from the uncertainties in long lived strange particle production and gluon splitting into heavy quark pairs. The effects of strange particle production are studied by varying the $s\bar{s}$ production probability in jet fragmentation. The $g \rightarrow b\bar{b}$ and $g \rightarrow c\bar{c}$ production rates are varied based upon the OPAL $g \rightarrow c\bar{c}$ measurement [10] and the theoretical prediction for the ratio $g \rightarrow b\bar{b}/g \rightarrow c\bar{c}$ [9].

The various charmed hadron production rates and fragmentation parameters in Z^0 decays are varied within the present LEP measurement errors. Charmed hadron fragmentation is studied by varying the average scaled energy $\langle x_E \rangle$ in the Peterson fragmentation function [11], as well as by studying the difference between the Peterson and Bowler models [12] for the same values of $\langle x_E \rangle$. Charmed hadron decay lifetimes are varied according to the world average measurement errors [13]. The charmed hadron decay charged multiplicity and K^0 production rate systematic uncertainties are based on measurements by Mark-III [14]. Charmed hadron decays with fewer neutral particles have higher charged mass and are therefore more likely to be tagged. Thus, an additional systematic uncertainty is estimated by varying the rates of charmed hadron decays with no π^0 s by $\pm 10\%$.

The B decay modeling uncertainty enters via the λ_b estimation. It is studied by varying the B lifetime, B baryon production rate, B fragmentation function and the B decay charged multiplicity in a similar manner as for the charm systematic studies. Simulation uncertainties which affect the tagging efficiency are studied by comparing the angular distribution of the b -tagging rate between data and simulation and a systematic error is assigned to the difference. Hard gluon radiation effects are estimated from $\pm 30\%$ variation of the fraction of simulation events where both B hadrons are contained within the same hemisphere and a hard gluon is in the other. Another systematic error is assigned to effects of B hadron momentum correlation between the two hemispheres, mostly due to soft gluon radiation and fragmentation effects, which in turn translate to a b -tagging efficiency correlation. This is estimated by comparing the B momentum correlation in the HERWIG [15] and JETSET [4] event generators.

As a cross check we decomposed the efficiency correlation into an independent set of components which represent all sources of correlation between the two b hemispheres. The components we have studied and their contributions are: the PV measurement (-0.02%), the track resolution effect on the IP determination ($+0.04\%$), the detector non-uniformity via the tagging angular distribution dependence ($+0.49\%$), the momentum distribution of the B hadron in each hemisphere ($+0.08\%$) and the effect of hard gluon emission forcing the two B hadrons into one hemisphere ($+0.07\%$). The estimated λ_b ($0.59 \pm 0.11\%$) and that from the sum of the components (0.67%) are in good agreement.

A major source of detector systematic uncertainty is due to the discrepancy in modeling the track impact parameter resolution, mainly along the beam axis. In the simulation track z impact parameters are smeared using a random Gaussian distribution of width $20 \mu\text{m} / \sin\theta$, as well as adjusted for z impact parameter mean position shifts to match the data. The full difference in R_b between the nominal and resolution-corrected samples is conservatively assigned to be the resolution systematic error. The difference between the measured and simulation charged track multiplicity as a function of $\cos\theta$ and momentum is attributed to an unsimulated tracking inefficiency correction. Both the tracking resolution and efficiency corrections require the use of a random number generator. After application of these corrections the result vary slightly with different random sequences. These fluctuations are included as an additional MC statistical uncertainty. The uncertainty on the primary vertex xy location simulation is estimated from the effect of adding a Gaussian tail to the IP distribution of $100 \mu\text{m}$ width for 0.5% of the simulated events. The simulation shows that the ≤ 3 jets requirement in the event selection favors $b\bar{b}$ over other $q\bar{q}$ events which biases our measurement by $+0.55\%$. We verified this bias in the data, by measuring R_b with and without applying the ≤ 3 jet criterion and found that our measured R_b value only changed by 0.0001, which is consistent with a statistical fluctuation but nevertheless was included as a systematic error. Another bias of $+0.26 \pm 0.12\%$ is introduced by the other event selection criteria, thus the combined bias is $+0.82 \pm 0.13\%$ and was corrected. Fig. 3(b) shows the statistical and the detector, physics and R_c systematic uncertainties versus the minimum \mathcal{M} cut.

In summary we have measured:

$$R_b = 0.2142 \pm 0.0034_{stat.} \pm 0.0015_{syst.} \pm 0.0002_{R_c}$$

which includes a correction of $+0.0003$ for the $e^+e^- \rightarrow \gamma \rightarrow b\bar{b}$ contribution. This value supersedes our previous R_b measurements [3] and is in good agreement with the SM prediction of 0.2155. A new high precision measurement has recently been reported by ALEPH [16], which also incorporates mass information to improve a lifetime-based probability tag. With the new SLD and LEP measurements the gap between the SM prediction of R_b and the world average has narrowed.

We thank the personnel of the SLAC accelerator department and the technical staffs of our collaborating institutions for their outstanding efforts on our behalf. This work was supported by the U.S. Department of Energy and National Science Foundation, the UK Particle Physics and Astronomy Research Council, the Istituto Nazionale di Fisica Nucleare of Italy, the Japan-US Cooperative Research Project on High Energy Physics, and the Korea Science and Engineering Foundation.

References

- [1] M. Hildreth proceedings of XXXI Rencontres de Moriond “*Electroweak Interactions and Unified Theories*”, J. Tran Than Van ed., (Editions Frontiers 1996) 147.

- [2] ALEPH Collab. D. Buskulic *et al.*, *Phys. Lett.* **B313** (1993) 535;
 OPAL Collab. P. D. Acton *et al.*, *Z. Phys.* **C60** (1993) 579; D. Akers
et al., *Z. Phys.* **C65** (1994) 17; K. Ackerstaff *et al.*, CERN-PPE-96-167,
 Submitted to *Z. Phys. C.* (1996);
 DELPHI Collab. P. Abreu *et al.*, *Z. Phys.* **C66** (1995) 323, *Z. Phys.* **C70**
 (1996) 531.
- [3] SLD Collab. K. Abe *et al.*, *Phys. Rev.* **D53**, (1996) 1023.
- [4] T. Sjostrand, *Comput. Phys. Commun.* **82** (1994) 74.
- [5] QQ MC code provided by P. Kim and the CLEO Collab. Tuning of the *B*
 decay modeling is described in SLD Collab., K. Abe *et al.*, *SLAC-PUB-*
7266 (1996), Submitted to *Phys. Rev. Lett.*
- [6] JADE Collab, S. Bethke *et al.*, *Phys. Lett.* **B213** (1988) 235.
- [7] D. Jackson, *Nucl. Inst. & Meth.* **A388** (1997) 247.
- [8] D. Bardin *et al.*, *CERN-TH 6443/92*, May 1992.
- [9] The LEP Electroweak Working Group, "Presentation of the LEP
 Electroweak Heavy Flavour Results for Summer 1996 Conferences",
LEPHF/96-01, July 1996.
- [10] OPAL Collab. R. Akers *et al.*, *Phys. Lett.* **B353** (1995) 595.
- [11] C. Peterson *et al.* *Phys. Rev.* **D27** (1983) 105.
- [12] M. G. Bowler, *Z. Phys.* **C11** (1981) 169.
- [13] Particle Data Group, *Phys. Rev.* **D50**, Part I (1994).
- [14] Mark-III Collab., D. Coffman *et al.*, *Phys Lett.* **B263** (1991) 135.
- [15] G. Marchesini *et al.*, *Comput. Phys. Commun.* **67** (1992) 465.
- [16] ALEPH Collab. R. Barate *et al.*, *Phys. Lett.* **B401** (1997) 150, 163.

*List of Authors

K. Abe,⁽¹⁹⁾ K. Abe,⁽³⁰⁾ T. Akagi,⁽²⁸⁾ N.J. Allen,⁽⁴⁾ W.W. Ash,^{(28)†} D. Aston,⁽²⁸⁾
K.G. Baird,⁽²⁴⁾ C. Baltay,⁽³⁴⁾ H.R. Band,⁽³³⁾ M.B. Barakat,⁽³⁴⁾ G. Baranko,⁽⁹⁾
O. Bardon,⁽¹⁵⁾ T. L. Barklow,⁽²⁸⁾ G.L. Bashindzhagyan,⁽¹⁸⁾ A.O. Bazarko,⁽¹⁰⁾
R. Ben-David,⁽³⁴⁾ A.C. Benvenuti,⁽²⁾ G.M. Bilei,⁽²²⁾ D. Bisello,⁽²¹⁾ G. Blaylock,⁽¹⁶⁾
J.R. Bogart,⁽²⁸⁾ B. Bolen,⁽¹⁷⁾ T. Bolton,⁽¹⁰⁾ G.R. Bower,⁽²⁸⁾ J.E. Brau,⁽²⁰⁾
M. Breidenbach,⁽²⁸⁾ W.M. Bugg,⁽²⁹⁾ D. Burke,⁽²⁸⁾ T.H. Burnett,⁽³²⁾ P.N. Burrows,⁽¹⁵⁾
W. Busza,⁽¹⁵⁾ A. Calcaterra,⁽¹²⁾ D.O. Caldwell,⁽⁵⁾ D. Calloway,⁽²⁸⁾ B. Camanzi,⁽¹¹⁾
M. Carpinelli,⁽²³⁾ R. Cassell,⁽²⁸⁾ R. Castaldi,^{(23)(a)} A. Castro,⁽²¹⁾ M. Cavalli-Sforza,⁽⁶⁾
A. Chou,⁽²⁸⁾ E. Church,⁽³²⁾ H.O. Cohn,⁽²⁹⁾ J.A. Coller,⁽³⁾ V. Cook,⁽³²⁾ R. Cotton,⁽⁴⁾
R.F. Cowan,⁽¹⁵⁾ D.G. Coyne,⁽⁶⁾ G. Crawford,⁽²⁸⁾ A. D'Oliveira,⁽⁷⁾ C.J.S. Damerell,⁽²⁵⁾
M. Daoudi,⁽²⁸⁾ N. de Groot,⁽²⁸⁾ R. De Sangro,⁽¹²⁾ R. Dell'Orso,⁽²³⁾ P.J. Dervan,⁽⁴⁾
M. Dima,⁽⁸⁾ D.N. Dong,⁽¹⁵⁾ P.Y.C. Du,⁽²⁹⁾ R. Dubois,⁽²⁸⁾ B.I. Eisenstein,⁽¹³⁾ R. Elia,⁽²⁸⁾
E. Etzion,⁽³³⁾ S. Fahey,⁽⁹⁾ D. Falciai,⁽²²⁾ C. Fan,⁽⁹⁾ J.P. Fernandez,⁽⁶⁾ M.J. Fero,⁽¹⁵⁾
R. Frey,⁽²⁰⁾ T. Gillman,⁽²⁵⁾ G. Gladding,⁽¹³⁾ S. Gonzalez,⁽¹⁵⁾ E.L. Hart,⁽²⁹⁾ J.L. Harton,⁽⁸⁾
A. Hasan,⁽⁴⁾ Y. Hasegawa,⁽³⁰⁾ K. Hasuko,⁽³⁰⁾ S. J. Hedges,⁽³⁾ S.S. Hertzbach,⁽¹⁶⁾
M.D. Hildreth,⁽²⁸⁾ J. Huber,⁽²⁰⁾ M.E. Huffer,⁽²⁸⁾ E.W. Hughes,⁽²⁸⁾ H. Hwang,⁽²⁰⁾
Y. Iwasaki,⁽³⁰⁾ D.J. Jackson,⁽²⁵⁾ P. Jacques,⁽²⁴⁾ J. A. Jaros,⁽²⁸⁾ Z. Y. Jiang,⁽²⁸⁾
A.S. Johnson,⁽³⁾ J.R. Johnson,⁽³³⁾ R.A. Johnson,⁽⁷⁾ T. Junk,⁽²⁸⁾ R. Kajikawa,⁽¹⁹⁾
M. Kalelkar,⁽²⁴⁾ H. J. Kang,⁽²⁶⁾ I. Karliner,⁽¹³⁾ H. Kawahara,⁽²⁸⁾ H.W. Kendall,⁽¹⁵⁾
Y. D. Kim,⁽²⁶⁾ M.E. King,⁽²⁸⁾ R. King,⁽²⁸⁾ R.R. Kofler,⁽¹⁶⁾ N.M. Krishna,⁽⁹⁾
R.S. Kroeger,⁽¹⁷⁾ J.F. Labs,⁽²⁸⁾ M. Langston,⁽²⁰⁾ A. Lath,⁽¹⁵⁾ J.A. Lauber,⁽⁹⁾
D.W.G.S. Leith,⁽²⁸⁾ V. Lia,⁽¹⁵⁾ M.X. Liu,⁽³⁴⁾ X. Liu,⁽⁶⁾ M. Loreti,⁽²¹⁾ A. Lu,⁽⁵⁾
H.L. Lynch,⁽²⁸⁾ J. Ma,⁽³²⁾ G. Mancinelli,⁽²⁴⁾ S. Manly,⁽³⁴⁾ G. Mantovani,⁽²²⁾
T.W. Markiewicz,⁽²⁸⁾ T. Maruyama,⁽²⁸⁾ H. Masuda,⁽²⁸⁾ E. Mazzucato,⁽¹¹⁾
A.K. McKemey,⁽⁴⁾ B.T. Meadows,⁽⁷⁾ R. Messner,⁽²⁸⁾ P.M. Mockett,⁽³²⁾ K.C. Moffeit,⁽²⁸⁾
T.B. Moore,⁽³⁴⁾ D. Muller,⁽²⁸⁾ T. Nagamine,⁽²⁸⁾ S. Narita,⁽³⁰⁾ U. Nauenberg,⁽⁹⁾ H. Neal,⁽²⁸⁾
M. Nussbaum,^{(7)†} Y. Ohnishi,⁽¹⁹⁾ N. Oishi,⁽¹⁹⁾ D. Onoprienko,⁽²⁹⁾ L.S. Osborne,⁽¹⁵⁾
R.S. Panvini,⁽³¹⁾ C.H. Park,⁽²⁷⁾ H. Park,⁽²⁰⁾ T.J. Pavel,⁽²⁸⁾ I. Peruzzi,^{(12)(b)} M. Piccolo,⁽¹²⁾
L. Piemontese,⁽¹¹⁾ E. Pieroni,⁽²³⁾ K.T. Pitts,⁽²⁰⁾ R.J. Plano,⁽²⁴⁾ R. Prepost,⁽³³⁾
C.Y. Prescott,⁽²⁸⁾ G.D. Punkar,⁽²⁸⁾ J. Quigley,⁽¹⁵⁾ B.N. Ratcliff,⁽²⁸⁾ T.W. Reeves,⁽³¹⁾
J. Reidy,⁽¹⁷⁾ P.L. Reinertsen,⁽⁶⁾ P.E. Rensing,⁽²⁸⁾ L.S. Rochester,⁽²⁸⁾ P.C. Rowson,⁽¹⁰⁾
J.J. Russell,⁽²⁸⁾ O.H. Saxton,⁽²⁸⁾ T. Schalk,⁽⁶⁾ R.H. Schindler,⁽²⁸⁾ B.A. Schumm,⁽⁶⁾
J. Schwiening,⁽²⁸⁾ S. Sen,⁽³⁴⁾ V.V. Serbo,⁽³³⁾ M.H. Shaevitz,⁽¹⁰⁾ J.T. Shank,⁽³⁾
G. Shapiro,⁽¹⁴⁾ D.J. Sherden,⁽²⁸⁾ K.D. Shmakov,⁽²⁹⁾ C. Simopoulos,⁽²⁸⁾ N.B. Sinev,⁽²⁰⁾
S.R. Smith,⁽²⁸⁾ M.B. Smy,⁽⁸⁾ J.A. Snyder,⁽³⁴⁾ H. Staengle,⁽⁸⁾ P. Stamer,⁽²⁴⁾ H. Steiner,⁽¹⁴⁾
R. Steiner,⁽¹⁾ M.G. Strauss,⁽¹⁶⁾ D. Su,⁽²⁸⁾ F. Suekane,⁽³⁰⁾ A. Sugiyama,⁽¹⁹⁾ S. Suzuki,⁽¹⁹⁾
M. Swartz,⁽²⁸⁾ A. Szumilo,⁽³²⁾ T. Takahashi,⁽²⁸⁾ F.E. Taylor,⁽¹⁵⁾ E. Torrence,⁽¹⁵⁾
A.I. Trandafir,⁽¹⁶⁾ J.D. Turk,⁽³⁴⁾ T. Usher,⁽²⁸⁾ J. Va'vra,⁽²⁸⁾ C. Vannini,⁽²³⁾ E. Vella,⁽²⁸⁾
J.P. Venuti,⁽³¹⁾ R. Verdier,⁽¹⁵⁾ P.G. Verdini,⁽²³⁾ D.L. Wagner,⁽⁹⁾ S.R. Wagner,⁽²⁸⁾
A.P. Waite,⁽²⁸⁾ S.J. Watts,⁽⁴⁾ A.W. Weidemann,⁽²⁹⁾ E.R. Weiss,⁽³²⁾ J.S. Whitaker,⁽³⁾
S.L. White,⁽²⁹⁾ F.J. Wickens,⁽²⁵⁾ D.C. Williams,⁽¹⁵⁾ S.H. Williams,⁽²⁸⁾ S. Willocq,⁽²⁸⁾

R.J. Wilson,⁽⁸⁾ W.J. Wisniewski,⁽²⁸⁾ M. Woods,⁽²⁸⁾ G.B. Word,⁽²⁴⁾ J. Wyss,⁽²¹⁾
R.K. Yamamoto,⁽¹⁵⁾ J.M. Yamartino,⁽¹⁵⁾ X. Yang,⁽²⁰⁾ J. Yashima,⁽³⁰⁾ S.J. Yellin,⁽⁵⁾
C.C. Young,⁽²⁸⁾ H. Yuta,⁽³⁰⁾ G. Zapalac,⁽³³⁾ R.W. Zdarko,⁽²⁸⁾ and J. Zhou,⁽²⁰⁾

- ⁽¹⁾ *Adelphi University, Garden City, New York 11530*
⁽²⁾ *INFN Sezione di Bologna, I-40126 Bologna, Italy*
⁽³⁾ *Boston University, Boston, Massachusetts 02215*
⁽⁴⁾ *Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom*
⁽⁵⁾ *University of California at Santa Barbara, Santa Barbara, California 93106*
⁽⁶⁾ *University of California at Santa Cruz, Santa Cruz, California 95064*
⁽⁷⁾ *University of Cincinnati, Cincinnati, Ohio 45221*
⁽⁸⁾ *Colorado State University, Fort Collins, Colorado 80523*
⁽⁹⁾ *University of Colorado, Boulder, Colorado 80309*
⁽¹⁰⁾ *Columbia University, New York, New York 10027*
⁽¹¹⁾ *INFN Sezione di Ferrara and Università di Ferrara, I-44100 Ferrara, Italy*
⁽¹²⁾ *INFN Lab. Nazionali di Frascati, I-00044 Frascati, Italy*
⁽¹³⁾ *University of Illinois, Urbana, Illinois 61801*
⁽¹⁴⁾ *E.O. Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720*
⁽¹⁵⁾ *Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*
⁽¹⁶⁾ *University of Massachusetts, Amherst, Massachusetts 01003*
⁽¹⁷⁾ *University of Mississippi, University, Mississippi 38677*
⁽¹⁸⁾ *Moscow State University, Institute of Nuclear Physics 119899 Moscow, Russia*
⁽¹⁹⁾ *Nagoya University, Chikusa-ku, Nagoya 464 Japan*
⁽²⁰⁾ *University of Oregon, Eugene, Oregon 97403*
⁽²¹⁾ *INFN Sezione di Padova and Università di Padova, I-35100 Padova, Italy*
⁽²²⁾ *INFN Sezione di Perugia and Università di Perugia, I-06100 Perugia, Italy*
⁽²³⁾ *INFN Sezione di Pisa and Università di Pisa, I-56100 Pisa, Italy*
⁽²⁴⁾ *Rutgers University, Piscataway, New Jersey 08855*
⁽²⁵⁾ *Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX United Kingdom*
⁽²⁶⁾ *Sogang University, Seoul, Korea*
⁽²⁷⁾ *Soongsil University, Seoul, Korea 156-743*
⁽²⁸⁾ *Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309*
⁽²⁹⁾ *University of Tennessee, Knoxville, Tennessee 37996*
⁽³⁰⁾ *Tohoku University, Sendai 980 Japan*
⁽³¹⁾ *Vanderbilt University, Nashville, Tennessee 37235*
⁽³²⁾ *University of Washington, Seattle, Washington 98195*
⁽³³⁾ *University of Wisconsin, Madison, Wisconsin 53706*
⁽³⁴⁾ *Yale University, New Haven, Connecticut 06511*

† *Deceased*

^(a) *Also at the Università di Genova*

^(b) *Also at the Università di Perugia*




Figure 1: (a) An illustration of the SV track attachment criteria. (b) Illustration of the P_t derivation.




Figure 2: Distribution of (a) M_{ch} and (b) P_t corrected mass, \mathcal{M} , for data (points) and MC which includes a breakdown of the b , c and uds contributions (open, hatched and crosshatched histograms respectively).




Figure 3: (a) Variation of R_b with the \mathcal{M} cut. The statistical error is represented by the boxes and the outer bars show the sum in quadrature of the statistical and systematic errors. (b) R_b statistical and systematic uncertainties versus \mathcal{M} cut.

Light Quark Systematic (ϵ_{uds})	δR_b
$g \rightarrow b\bar{b}$ $0.31 \pm 0.11\%$	-0.00033
$g \rightarrow c\bar{c}$ $2.38 \pm 0.48\%$	-0.00004
K^0 production $\pm 10\%$	-0.00003
Λ production $\pm 10\%$	-0.00002
Total uds physics systematic	0.00034
Charm Systematic (ϵ_c)	δR_b
D^+ production 0.259 ± 0.028	-0.00011
D_s production 0.115 ± 0.037	-0.00005
c -baryon production 0.074 ± 0.029	0.00011
c -frag. $\langle x_E \rangle_D = 0.482 \pm 0.008$	-0.00006
c -frag. function shape	-0.00001
D^0 lifetime 0.415 ± 0.004 ps	-0.00003
D^+ lifetime 1.057 ± 0.015 ps	-0.00001
D_s lifetime 0.467 ± 0.017 ps	-0.00002
Λ_c lifetime 0.200 ± 0.011 ps	-0.00001
D^0 decay $\langle N_{ch} \rangle = 2.54 \pm 0.05$	-0.00006
D^+ decay $\langle N_{ch} \rangle = 2.50 \pm 0.06$	-0.00006
D_s decay $\langle N_{ch} \rangle = 2.65 \pm 0.33$	-0.00009
$D^0 \rightarrow K^0$ production 0.401 ± 0.059	+0.00015
$D^+ \rightarrow K^0$ production 0.646 ± 0.078	+0.00020
$D_s \rightarrow K^0$ production 0.380 ± 0.06	+0.00002
D^0 decay no- π^0 frac. 0.370 ± 0.037	+0.00005
D^+ decay no- π^0 frac. 0.499 ± 0.050	-0.00008
D_s decay no- π^0 frac. 0.352 ± 0.035	<0.00001
Total Charm Physics systematic	0.00033
B decay modeling (λ_b)	δR_b
B lifetime ± 0.05 ps	0.00004
B decay $\langle N_{ch} \rangle = 5.73 \pm 0.35$	0.00003
b fragmentation	0.00019
Λ_b production fraction 0.074 ± 0.03	0.00008
Hard gluon radiation	0.00008
B momentum correlation	0.00029
b -tag $\cos \theta$ dependency	0.00001
Total $b\bar{b}$ Physics systematic	0.00038
Detector Systematic	δR_b
Tracking resolution	0.00096
Tracking efficiency	0.00040
$\langle IP \rangle_{xy}$ tail	0.00010
MC statistics	0.00091
Event selection bias	0.00028
Total detector and MC	0.00141
$R_c = 0.171 \pm 0.006$	0.00021
Total (excl. R_c)	0.00154

Table 1: Summary of systematic uncertainties for the $\mathcal{M} > 2.0$ GeV/ c^2 cut.