

Singapore Management University

Institutional Knowledge at Singapore Management University

Research Collection School of Social Sciences

School of Social Sciences

1-2023

COVID-19 stress and cognitive failures in daily life: A multilevel examination of within- and between-persons patterns

NADYANNA BINTE MOHAMED MAJEED

Singapore Management University, nadyannam.2020@msps.smu.edu.sg

K TENNAKON APPUHAMILLAGA SANDEESHWARA KASTURIRATNA

Singapore Management University, skasturirat.2023@phdps.smu.edu.sg

Ming Yao LI

Jonathan L. CHIA

Verity Y. Q. LUA

See next page for additional authors

Follow this and additional works at: https://ink.library.smu.edu.sg/sooss_research



Part of the [Asian Studies Commons](#), [Cognitive Psychology Commons](#), [Public Health Commons](#), and the [Social Psychology Commons](#)

Citation

NADYANNA BINTE MOHAMED MAJEED, , K TENNAKON APPUHAMILLAGA SANDEESHWARA KASTURIRATNA, , LI, Ming Yao, CHIA, Jonathan L., LUA, Verity Y. Q., & HARTANTO, Andree.(2023). COVID-19 stress and cognitive failures in daily life: A multilevel examination of within- and between-persons patterns. *Journal of Pacific Rim Psychology, 17*, 1-10.
Available at: https://ink.library.smu.edu.sg/sooss_research/3839

This Journal Article is brought to you for free and open access by the School of Social Sciences at Institutional Knowledge at Singapore Management University. It has been accepted for inclusion in Research Collection School of Social Sciences by an authorized administrator of Institutional Knowledge at Singapore Management University. For more information, please email cherylids@smu.edu.sg.

Author

NADYANNA BINTE MOHAMED MAJEED, K TENNAKOON APPUHAMILLAGE SANDEESHWARA
KASTURIRATNA, Ming Yao LI, Jonathan L. CHIA, Verity Y. Q. LUA, and Andree HARTANTO

COVID-19 stress and cognitive failures in daily life: A multilevel examination of within- and between-persons patterns

Journal of Pacific Rim Psychology

Volume 17: 1–10

© The Author(s) 2023

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/18344909231208119

journals.sagepub.com/home/pac



Nadyanna M. Majeed^{1,2,*} , K. T. A. Sandeeshwara Kasturiratna^{2,*} ,
Ming Yao Li² , Jonathan L. Chia² , Verity Y. Q. Lua^{2,3}
and Andree Hartanto²

Abstract

The COVID-19 pandemic has claimed an extremely high number of lives worldwide, causing widespread panic and stress. The current research examined whether COVID-19 stress was associated with everyday cognitive failures, using data from a seven-day daily diary study of 253 young adults in Singapore. Multilevel modeling revealed that COVID-19 stress was significantly associated with cognitive failures even after adjusting for demographic factors, both at the within-person and between-persons levels. Specifically, individuals experienced more cognitive failures on days they experienced more COVID-19 stress (as compared to their own average levels of COVID-19 stress), and individuals who experienced more COVID-19 stress overall (as compared to individuals who experienced less COVID-19 stress overall) experienced more cognitive failures in general. While a large body of work has evidenced the detrimental effects of COVID-19 stress on individuals' well-being, the current findings provide novel insights that these stressors may negatively impact individuals' cognitive functioning as well.

Keywords

COVID-19, stress, cognitive failures, daily diary, multilevel modeling

Received 8 May 2022; accepted 23 September 2023

COVID-19 stress and cognitive failures in daily life

The COVID-19 pandemic has claimed an extremely high number of lives worldwide. As of early April 2022, there have been more than 505 million confirmed cases of COVID-19 infection and more than 6 million deaths (World Health Organization, 2022). Beyond the loss of life, the fear of potential infection and constant worry about the health of loved ones have resulted in significant stress and anxiety for many individuals (Xu et al., 2020). Moreover, the resultant abrupt changes from higher unemployment rates and social isolation have caused widespread financial insecurity and loneliness, compounding the experienced stress during the pandemic (Vatansever et al., 2021). Furthermore, unlike other daily stressors that occur due to various context-dependent factors specific to each person (e.g., relationship stressors, work-related stressors), COVID-19 related stressors can be experienced by almost everyone concurrently in the world today (Arıcan & Kafadar, 2022). Thus, COVID-19 related stressors are a

more widespread phenomenon in the current background that is particularly important to study. While the economic and socioemotional consequences of COVID-19 stress have been well established, little is currently known about its cognitive consequences.

¹National University of Singapore, Singapore, Singapore

²Singapore Management University, Singapore, Singapore

³Stanford University, Stanford, California

*NMM and KTASK should be considered joint first author. The order of co-first authors provided here was randomised. Co-first authors can prioritise their names when adding this paper's reference to their resumes.

Corresponding authors:

Nadyanna M. Majeed, Faculty of Arts & Social Sciences, National University of Singapore, 5 Arts Link, Level 5, Singapore 117570, Singapore. Email: nadyanna@u.nus.edu

Andree Hartanto, School of Social Sciences, Singapore Management University, 10 Canning Rise, Level 5, Singapore 179873, Singapore. Email: andreeh@smu.edu.sg



One potential cognitive consequence of COVID-19 stress is related to daily cognitive failures. Cognitive failures are broadly defined as failures to perform simple tasks that a person is normally capable of completing without errors (Lange & Süß, 2014), such as stopping at the traffic light when it turns red or remembering to attend a work meeting. Cognitive failures can be distinguished from other cognitive errors as being the result of lapses of attention or slips of memory rather than a result of a lack of knowledge, and individuals who experience cognitive failure more often report frequent lapses of action, perception, and memory (Bridger et al., 2013). Although cognitive slips and errors are common in daily living, they have been found to have broad and potentially major implications in real-world outcomes. For example, cognitive failures make individuals more vulnerable to the effects of stress (Broadbent et al., 1982; Mahoney et al., 1998), leading to outcomes such as job impairment, workplace-related mishaps, and injuries and hospitalization (Abbasi et al., 2021; Wallace & Vodanovich, 2003a, 2003b). Cognitive failures have also been found to lead to increased susceptibility to driving error and car accidents (Allahyari et al., 2019; Larson & Merritt, 1991). Several studies also indicate that self-reported cognitive failures predict laboratory measures assessing cognitive abilities such as working memory or attentional control (Unsworth et al., 2012a; Unsworth et al., 2012b).

Research examining the antecedents of cognitive failures has suggested that stress can have detrimental effects on one's cognitive functioning (Mahoney et al., 1998). In particular, it is thought that the negative effect of stress on cognitive functioning may be explained through the mediating role of mind wandering (Hartanto & Yang, 2020), which refers to thoughts about one's goals that are not related to the primary task at hand (Morin & Racy, 2021; Smallwood & Schooler, 2006). While mind-wandering is not necessarily maladaptive, it can negatively affect cognitive functioning if driven by anxiety or stress, and lead to cognitive failures (Burdett et al., 2016; McVay & Kane, 2010). For example, stressful life events, such as the COVID-19 pandemic, may increase the likelihood of engaging in task-unrelated thoughts (e.g., worrying about potential infection) and decrease the effectiveness of disengaging from such thoughts (Klein & Boals, 2001). These mental diversions compete for limited cognitive resources and the attempts to suppress undesired thoughts consume even more resources (Smeekens & Kane, 2016). Indeed, as illustrated by literature on working memory, these mental diversions may result in poorer reasoning (Kane et al., 2005), multitasking (Colom et al., 2010), and learning (Noël, 2009). Given the literature on the implications of cognitive failure, as well as the pervasiveness of stress during this pandemic, it behooves us to properly examine the relationship between COVID-19 stress and cognitive failure.

The present study thus seeks to investigate the effect of COVID-19 stress on everyday cognitive failures. In order to reduce the possibility that individual differences may

confound any observed associations between COVID-19 stress and cognitive failures, we employed a multilevel approach by utilizing a diary study design. This study design allowed us to examine day-to-day fluctuations in COVID-19 stress and cognitive failures, while controlling for several individual differences in the form of demographic factors (i.e., age, sex, race, household income, and subjective socioeconomic status) previously linked to our variables of interest.

Method

Participants and design

Data for the current study were collected as part of the second wave of a larger project broadly examining daily experiences and well-being (see Goh et al., 2023; Majeed et al., 2023; Ng, Lua et al., 2022; Ng, Majeed et al., 2023), conducted with young adults in Singapore with the daily diary portions running from June 24, 2021 to August 12, 2021 (i.e., during the country's partially relaxed "Phase 3 Heightened Alert"¹ and the stricter semi-lockdown "Phase 2 Heightened Alert"² of the COVID-19 pandemic; Abu Baker, 2021; Ang, 2021).

Undergraduate students from the last author's university were recruited for the study through the local subject pool system.³ Participants completed a baseline questionnaire, followed by a seven-day self-administered diary survey taken nightly. Within one to two days of completing the baseline questionnaire, a link to the first diary survey was emailed to participants at 8.00 p.m., with the link to the consecutive diary surveys being sent each evening at 8.00 p.m. for the remaining six days. Each daily diary survey was left available for completion until 3.00 a.m. the next day. A total of 1,721 observations of daily data were obtained from 253 participants, where each participant provided an average of 6.80 days of observations (97% completion rate; see Table 1). The data collection procedure was approved by the Institutional Review Board at the last author's university (IRB-20-118-A075-M4[621]). All participants provided informed consent prior to data collection.

Measures

Baseline. Demographic covariates. Participants were asked to report their sex (male or female), race (Chinese, Malay, Indian, or other), age, and monthly household income (1 = less than \$2,000, 2 = \$2,000–\$5,999, 3 = \$6,000–\$9,999, 4 = \$10,000–\$14,999, 5 = \$15,000–\$19,999, 6 = more than \$20,000). Subjective socioeconomic status was measured using a 10-point ladder scale (Adler et al., 2000), with the bottom rung (1) representing the lowest status and the top rung (10) representing the highest status.

COVID-19 vaccination status. Participants were asked to report their COVID-19 vaccination status (currently vaccinated or unvaccinated).

Table 1. Summary of descriptive statistics

Variable	N	M	SD	Observed Range	Theoretical Range
Person level					
Sex (% female)	253	77%			
Race (% Chinese)	253	75%			
Age (in years)	253	22.11	1.63	19–29	
Monthly household income	253	3.00	1.43	1–6	1–6
Subjective socioeconomic status	253	6.11	1.25	2–10	1–10
Vaccination status (% vaccinated)	253	71%			
Average stressor exposure	253	.37	.29	.00–1.00	.00–1.00
Average COVID-19 stress	253	0.32	0.46	0.00–2.63	0.00–4.00
Average cognitive failures	253	0.34	0.35	0.00–2.87	0.00–3.00
Day level					
Daily stressor exposure (% stressor days)	1721	38%			
Daily COVID-19 stress	1721	0.32	0.52	0.00–3.60	0.00–4.00
Daily cognitive failures	1721	0.33	0.44	0.00–3.00	0.00–3.00

Note: Ns for person level refer to number of participants, while Ns for day level refer to number of observations (days).




Figure 1. Participant-average levels of COVID-19 stress and national numbers of COVID-19 cases and deaths over time.

Note: Time is indicated in terms of days since the start of the daily diary portion of data collection for the current work. That is, the first day of response is day 1.

Daily. Daily stressor exposure. Daily stressor exposure was measured each day for each participant through the Daily Inventory of Stressful Events (Almeida et al., 2002). The inventory is a checklist of seven types of stressors (arguments, avoided arguments, discrimination, work/education stressors, home stressors, network stressors, and “others”), and participants were asked to indicate if any of the stressors occurred to them “since yesterday.” In line with existing work on daily stressor exposure (e.g., Majeed et al., 2021; Ng et al., 2022; Rush et al., 2019), the construct was operationalized as a binary variable. Specifically, if at least one stressor was experienced on that day, the day was categorized as a stressor day, and otherwise, it was categorized as a

non-stressor day. The intra-class correlation for daily stressor exposure was .64.

Daily COVID-19 stress. COVID-19 stress was measured using five items selected from the COVID Stress Scales (Taylor et al., 2020), administered daily. The COVID Stress Scales have been validated across a number of different samples (e.g., Asmundson et al., 2022; Carlander et al., 2022; Khosravani et al., 2021; Taylor et al., 2020), and correlate strongly with other COVID-19 fear-related measures (e.g., Khosravani et al., 2021; Pakpour et al., 2021) while showing significantly greater correlations with anxiety-related measures rather than depression-related measures (Khosravani et al., 2021; Taylor et al., 2020). The current study used five items adapted from the COVID Stress Scales to measure COVID-19-related stress and anxiety symptoms on four factors: (1) danger and contamination, (2) xenophobia, (3) compulsive checking, and (4) traumatic stress symptoms. Participants were asked to indicate “how you have felt or behaved this way today” in relation to three worries and two behaviors on a 5-point Likert scale (0 = *Not at all*, 4 = *Extremely* for worries; 0 = *Never*, 4 = *Almost always* for behaviors). Specifically, the items on worries were: “Today, I was worried about catching the virus” (danger and contamination), “Today, I was worried that our healthcare system will be unable to keep me and my loved ones safe from the virus” (danger and contamination), and “Today, I was worried that foreigners are spreading the virus in Singapore” (xenophobia). The items on behaviors were: “Today, I checked online and offline media (e.g., Facebook, Straits Times) for information concerning COVID-19” (compulsive checking) and “Today, I had trouble concentrating because I kept thinking about the virus” (traumatic stress symptoms). All items were averaged such that higher overall scores reflected greater levels of daily COVID-19 stress ($\alpha_{\text{within}} = .63$, $\alpha_{\text{between}} = .86$). The intra-class correlation for daily COVID-19 stress was .74. A plot




Figure 2. Conceptual representation of mediation.

of average COVID-19 stress levels (per participant) over time, in comparison to the number of COVID-19 cases and deaths in Singapore (data retrieved from Mathieu et al., 2020), is provided in Figure 1.

Daily cognitive failures. Cognitive failure was measured using the 13-item Cognitive Failures in Everyday Life Scale (Lange & Süß, 2014), administered daily. Each item (e.g., “Did your mind unintentionally wander, at any point of time today?”) was rated on a 4-point Likert scale (0 = *Never*, 3 = *Several times*), with higher scores reflecting greater daily cognitive failure ($\alpha_{\text{within}} = .75$, $\alpha_{\text{between}} = .95$). The intra-class correlation for daily cognitive failures was .57.

Analytic plan

Due to the two-level nature of the diary data (i.e., repeated measures nested within individuals), multilevel modeling⁴ was conducted to examine the association between daily COVID-19 stress and daily cognitive failures across the seven days (Level 1) nested within participants (Level 2).

Main analyses. The Level 1 predictor, daily COVID-19 stress, was person-mean centered and entered as both a random and fixed predictor, and its person-level mean was reintroduced as a fixed predictor at Level 2 in order to fully decompose daily COVID-19 stress into its within- and between-persons components (Bolger et al., 2012). Time—in terms of number of days since the start of data collection for the whole sample—was included as a person-mean-centered Level 1 covariate with both random and fixed components with its person-level mean reintroduced as a Level 2-fixed covariate in order to account for changes over time due to the changing pandemic situation. The unadjusted model was thus as follows, with γ_{10} and γ_{01} representing the within-persons and between-persons parameters of interest respectively:

Level 1: (Daily cognitive failures)_{di} = B_{0i} + B_{1i} (daily COVID-19 stress)_{di} + B_{2i} (time)_{di} + ϵ_{di}

Level 2: B_{0i} = γ_{00} + γ_{01} (average COVID-19 stress)_i + γ_{02} (average time)_i + μ_{0i}

B_{1i} = γ_{10} + μ_{1i}

B_{2i} = γ_{20} + μ_{2i}

An adjusted model was estimated with the inclusion of demographic factors, COVID-19 vaccination status, and daily stressor exposure. Age, sex, race, objective socioeconomic status (i.e., monthly household income), and

subjective socioeconomic status were included as Level 2-fixed components in an adjusted model in order to rule out demographic confounds. Stressor exposure was included as random and fixed components at Level 1 in terms of stressor days versus non-stressor days, and as a fixed component at Level 2 in terms of average stressor exposure across the seven days. Categorical predictors (i.e., sex, race, and daily stressor exposure) were dummy-coded with *male*, *majority race* (i.e., *Chinese*), and *non-stressor day* serving as the reference categories respectively. The equations for the adjusted model are thus as follows:

Level 1: (Daily cognitive failures)_{di} = B_{0i} + B_{1i} (daily COVID-19 stress)_{di} + B_{2i} (time)_{di} + B_{3i} (daily stressor exposure)_{di} + ϵ_{di}

Level 2: B_{0i} = γ_{00} + γ_{01} (average COVID-19 stress)_i + γ_{02} (average time)_i + γ_{03} (age)_i + γ_{04} (sex)_i + γ_{05} (race)_i + γ_{06} (objective SES)_i + γ_{07} (subjective SES)_i + γ_{08} (average stressor exposure)_i + μ_{0i}

B_{1i} = γ_{10} + μ_{1i}

B_{2i} = γ_{20} + μ_{2i}

B_{3i} = γ_{30} + μ_{3i}

At Level 1, B_{0i} (the intercept coefficient for each participant *i*) indicates participant *i*'s average level of cognitive failures. B_{1i} shows the change in daily cognitive failures in relation to daily COVID-19 stress. At Level 2, B_{0i} was modeled as a function of between-participant differences. This includes participant *i*'s average COVID-19 stress over the duration of the seven days and other covariates. B_{1i} represents the slope coefficient for each participant *i*. μ_{0i} and μ_{1i} indicate the deviation of each participant's intercept and slope respectively from the model-implied values.

Additional analyses. Leave-one-out multiverse analysis. As the daily COVID-19 Stress measure was adapted and had not yet been established in its current form, we conducted a leave-one-out multiverse analysis to evaluate plausible analytic alternatives in parallel and interpreted collectively. The purpose for this analysis was to see if our results were robust across the different operationalizations of COVID-19 stress. Specifically, we repeated the unadjusted and adjusted analyses previously mentioned but with all 31 possible combinations of the five items used to create the COVID-19 Stress measure.

Exploratory lagged analysis. We repeated the main analyses with a one-day time lag in a preliminary examination of whether the association between COVID-19 stress and cognitive failures could spill over onto the next day. Specifically, we modeled next-day cognitive failures as a function of today's COVID-19 stress and covariates.

Exploratory mediation analysis. We repeated the main analyses with COVID-19 vaccination status as a predictor

of daily COVID-19 stress, resulting in a mediation model (for conceptual representation, see Figure 2).

Transparency and openness

The current work's design and its analysis plan were not pre-registered. Relevant materials, data, analytic code, as well as a supplemental information (zero-order correlation matrix at Level 2 and residual QQ plot) have been made publicly available on Researchbox #466 (<https://researchbox.org/466>).

Unless otherwise stated, all analyses were conducted in R version 3.6.3 (R Core Team, 2020). Data cleaning and manipulation were performed using *dplyr* version 1.0.10 (Wickham et al., 2022). Descriptives were extracted using *psych* version 2.2.9 (Revelle, 2021). Multilevel scale reliabilities were calculated via *semTools* version 0.5-5 (Jorgensen et al., 2021). ICCs were calculated by *merTools* version 0.5.2 (Knowles et al., 2020). Multilevel modeling was conducted via *lme4* version 1.1-28 (Bates et al., 2014) with significance testing by *lmerTest* version 3.1-3 (Kuznetsova et al., 2017), where maximum likelihood estimation was used to fit all models. Effect sizes in the form of standardized coefficients were calculated by *effectsize* version 0.6.0.1 (Ben-Shachar et al., 2020) using the "pseudo" method. Multilevel mediation was conducted via *mlma* version 6.2-1 (Yu & Li, 2022). Sensitivity power analyses were performed using *Mplus* version 8 (Muthén & Muthén, 2012).

Results

Main analyses

We found significant positive associations between daily COVID-19 stress and daily cognitive failures both at the within-persons and between-persons levels (Table 2).⁵ Specifically, to a small extent, participants reported more cognitive failures on days they experienced more COVID-19 stress compared to their own average (unadjusted: $\beta = .07$, 95% CI = [.02, .12], $\gamma_{10} = 0.08$, $SE = 0.03$, $p = .006$; adjusted: $\beta = .06$, 95% CI = [.01, .11], $\gamma_{10} = 0.07$, $SE = 0.03$, $p = .020$). Additionally, to a moderate extent, participants who experienced higher levels of COVID-19 stress in general also reported more cognitive failures than participants who experienced lower levels of COVID-19 stress (unadjusted: $\beta = .17$, 95% CI = [.04, .30], $\gamma_{01} = 0.12$, $SE = 0.05$, $p = .009$; adjusted: $\beta = .18$, 95% CI = [.09, .27], $\gamma_{01} = 0.13$, $SE = 0.03$, $p < .001$).

Leave-one-out multiverse analyses

Leave-one-out multiverse analyses revealed that our main findings were generally robust across the various operationalizations of daily COVID-19 stress (Figure 3). All point estimates were consistent with the main findings (i.e., positive in sign), although some combinations of items (e.g., the combination of items 1, 2, and 4 at the within-persons level,

Table 2. Full results of main multilevel analyses

Predictor	Unadjusted				Adjusted			
	Std. Est.	Est.	SE	<i>p</i>	Std. Est.	Est.	SE	<i>p</i>
Fixed effects								
Intercept	.00	0.39	0.07	<.001	.00	0.18	0.27	.509
<i>Within-persons</i>								
Time	-.12	-0.02	0.00	<.001	-.10	-0.01	0.00	.001
COVID-19 stress	.07	0.08	0.03	.006	.06	0.07	0.03	.019
Stressor exposure					.20	0.11	0.02	<.001
<i>Between-persons</i>								
Time	-.10	0.00	0.00	.108	-.05	0.00	0.00	.281
COVID-19 stress	.17	0.12	0.05	.009	.18	0.13	0.03	<.001
Stressor exposure					.41	0.46	0.06	<.001
Age					-.01	0.00	0.01	.817
Sex (0 = male, 1 = female)					.09	0.07	0.04	.094
Race (0 = majority, 1 = minority)					.01	0.00	0.04	.892
Objective SES					-.02	-0.01	0.01	.641
Subjective SES					-.02	-0.01	0.01	.626
Random effects								
Intercept		0.11				0.04		
Time		0.00				0.00		
COVID-19 stress		0.01				0.01		
Stressor exposure						0.03		
Residual		0.07				0.07		

Note: $N_{\text{participants}} = 253$, $N_{\text{observations}} = 1721$. Estimates for random effects indicate variances. Bolded rows indicate statistical significance at $p < .050$.

and the combination of items 1 and 4 at the between-persons level) produced nonsignificant associations ($p \geq .050$) with daily cognitive failures.

Exploratory lagged analyses

We found that daily COVID-19 stress was not predictive of next-day cognitive failures (unadjusted: $\beta = -.02$, 95% CI = $[-.07, .04]$, $\gamma_{10} = -0.02$, $SE = 0.03$, $p = .525$; adjusted: $\beta = -.02$, 95% CI = $[-.07, .03]$, $\gamma_{10} = -0.02$, $SE = 0.03$, $p = .455$). This suggests that the association between

COVID-19 stress and cognitive failures is short-lived, and does not spill over to the next day.

Exploratory mediation analyses

The mediation analysis revealed a significant indirect mediation pathway from COVID-19 vaccination status through COVID-19 stress in predicting cognitive failures ($B = -.01$, 95% CI = $[-.007, -.017]$). This suggests that being vaccinated for COVID-19 is associated with




Figure 3. Summary of leave-one-out multiverse analyses.

Note: Circles represent unstandardized point estimates while horizontal lines represent corresponding 95% CIs. Green indicates statistical significance at $p < .050$, red indicates non-significance, and grey indicates potentially inaccurate estimates due to singularity issues during the model fitting process. Black dashed vertical line indicates the value of the point estimate for the overall 5-item measure.

decreased levels of COVID-19 stress, which in turn is associated with lower levels of cognitive failures.

Discussion

We analyzed data from a daily diary study of 253 young adults in Singapore across seven days during the COVID-19 pandemic. We found that COVID-19 stress was consistently associated with poorer cognitive functioning both at the within-persons and between-persons levels, such that individuals experienced more cognitive failures on days they experienced more COVID-19 stress (as compared to their own average levels of COVID-19 stress), and also that individuals who experienced more COVID-19 stress overall (as compared to individuals who experienced less COVID-19 stress overall) experienced more cognitive failures in general. Adjusting for demographic covariates did not change the observed patterns. Of note, leave-one-out multivariate analyses of the five items forming the COVID-19 stress measure preliminarily suggest that the associations observed in the current work are mostly driven by stress from two specific items (i.e., “Today, I was worried that foreigners are spreading the virus in Singapore” and “Today, I had trouble concentrating because I kept thinking about the virus”).

The current work is the first to empirically examine the associations between COVID-19 stress and cognitive failures. The use of a daily diary methodology allowed us to examine this relationship in an ecologically valid setting, and also allowed us to rule out stable individual differences that may account for the positive within-persons associations between COVID-19 stress and cognitive failure. Taken together, the findings of the current work suggest that the stress arising from the COVID-19 situation may indeed be associated with increased levels of cognitive failures among individuals both at the day-to-day level and in general. The positive associations observed in the current study are in line with much of the literature broadly examining stress on cognitive failures (Mahoney et al., 1998). Considering past works suggesting that mind-wandering mediates the positive relationship between stress and cognitive failures (Hartanto & Yang, 2020), we posit that mind-wandering may also be a possible explanation for the observed relationship between COVID-19 stress and cognitive failure. Further research into this potential mechanism is warranted, given the lack of direct examination of this mechanism in the current work. Nonetheless, our findings suggest that beyond detriments to affective and socio-economic outcomes, COVID-19 stress may be associated with poorer cognitive outcomes as well.

In addition to our main results, we uncovered two notable exploratory findings. First, we did not observe any lagged associations or spillover of daily COVID-19 stress onto cognitive failures on the next day. This contrasts

with existing works on other types of stress and cognitive failure—for example, daily work-related stress has been found to be associated with higher levels of next-day cognitive failure (Kottwitz et al., 2022). Situating our current exploratory findings within the existing literature suggests that the relationship between COVID-19 stress and cognitive failure may be more transient in nature, perhaps due to differences in the nature of the stressor experience itself. Second, vaccination against COVID-19 appeared to play a protective role, with our current data indicating that vaccinated individuals reported reduced levels of COVID-19 stress in general. Consequently, this reduction in stress might indirectly protect against cognitive failures. These exploratory findings, while brief, provide valuable insights that could guide future research and interventions not just within the context of COVID-19, but within the broader stress literature.

The current study is not without limitations. First, while the original version of the COVID-19 Stress Scale has been validated with two large North American samples (Taylor et al., 2020) and a large South American sample (Pulido & Jimenez, 2020), it has not been validated with any Asian samples, nor has the shortened daily version adapted for the current daily diary study been validated in any context. As such, future studies should be conducted to assess the psychometric properties of the five-item version employed in the current study. Second, while the daily diary method is high in ecological validity, minimizes issues of recall bias, and allows for the disentangling of within- and between-persons associations, its nonexperimental nature does not allow for causal conclusions to be drawn. Thus, it is possible—though not theoretically likely—that higher levels of cognitive failures drive higher levels of COVID-19 stress,⁶ in opposition to our hypothesized direction.

In sum, the present work provides novel insights into the associations between daily COVID-19 stress and daily cognitive failures. While a large body of work has evidenced the detrimental effects of COVID-19 stress on individuals’ sense of well-being, the current findings suggest that these stressors may negatively impact individuals’ cognitive functioning as well. In turn, this may have unfavorable consequences on individuals’ day-to-day functioning (e.g., in logical reasoning and decision-making), as previously alluded to. Considering the pervasiveness of COVID-19 stress today and the high likelihood that the pandemic will continue affecting individuals’ lives in the near future (Chia & Hartanto, 2021), it is undoubtedly important to consider the downstream effects of the pandemic on individuals in the society. Beyond ensuring adequate medical support and infrastructure in the current pandemic, it is crucial for community leaders and policymakers to ensure that individuals have adequate resources to mitigate the stresses arising from the pandemic situation.

Acknowledgements

We are grateful to Frosch Quek, Gloria Lai, Harkiran Kaur, Harleen Kaur, Joax Wong, Manmeet Kaur, Wei Ming Ooi, and Xin Yi Poh for their assistance in data collection.

Authors contributions

NMM and AH developed the study concept and analytic plan. NMM and KTASK performed the data analysis and interpretation. NMM, KTASK, MYL, JLC, and VYQL drafted the paper, and all authors provided critical revisions. All authors approved the final version of the paper for submission.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding


The author(s) received no financial support for the research, authorship, and/or publication of this article: This research was supported by a grant awarded to AH by Singapore Management University through research grants from the Ministry of Education Academy Research Fund Tier 1 (21-SOSS-SMU-023) and Lee Kong Chian Fund for Research Excellence. The funding body had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Data availability statement

The materials and dataset analyzed during the current study have been made publicly available in the Researchbox repository (#466; <https://researchbox.org/466>).

ORCID iDs

Nadyanna M. Majeed  <https://orcid.org/0000-0003-0148-0846>

K. T. A. Sandeeshwara Kasturiratna  <https://orcid.org/0000-0002-7339-3875>

Jonathan L. Chia  <https://orcid.org/0000-0002-1055-5056>

Notes

1. Phase 3 Heightened Alert involved anti-pandemic measures such as limiting social gatherings to five people, restrictions on dining-in in eateries, restrictions on social rituals such as wedding receptions.
2. Phase 2 Heightened Alert was a stricter version of Phase 3 Heightened alert and involved semi-lockdown measures such as limiting the number of household visitors to two per day, enforcing work-from-home as the default mode of employment, and disallowing dining-in at eateries.
3. Only participants who did not participate in the first wave of the larger project broadly examining daily experiences and well-being were recruited for the study.
4. Visual observation of the residuals (plots available in Researchbox #466) suggested that the residuals were not normally distributed, potentially violating the assumption of normality. However, simulations by Maas and Hox (2004) showed that the estimates of interest derived from multilevel

modeling are generally robust to the normality assumption violation.

5. Sensitivity power analyses for multilevel designs (see Bolger et al., 2012) revealed that at an alpha level of .05, 80% power, and the current study's sample size of 253 participants and a maximum of 1,771 completed daily interviews, we would be able to detect a minimum γ_{10} of 0.07855, and a minimum γ_{01} of 0.08527.
6. We found that with cognitive failures as the predictor and COVID-19 stress as the outcome, there were significant positive associations between daily cognitive failures and daily COVID-19 stress both at the within-persons (unadjusted: $\beta = .08$, 95% CI = [.02, .14], $\gamma_{10} = 0.40$, $SE = 0.14$, $p = .008$; adjusted: $\beta = .08$, 95% CI = [.02, .13], $\gamma_{10} = 0.40$, $SE = 0.14$, $p = .008$) and between-persons (unadjusted: $\beta = .18$, 95% CI = [.06, .31], $\gamma_{01} = 1.17$, $SE = 0.40$, $p = .004$; adjusted: $\beta = .20$, 95% CI = [.06, .34], $\gamma_{01} = 1.28$, $SE = 0.46$, $p = .006$) levels. Nevertheless, we caution that this direction of the association (i.e., cognitive failure \rightarrow COVID-19 stress) is not theoretically sound.

References

- Abbasi, M., Falahati, M., Kaydani, M., Fallah Madvari, R., Mehri, A., Ghaljahi, M., & Yazdanirad, S. (2021). The effects of psychological risk factors at work on cognitive failures through the accident proneness. *BMC Psychology*, 9(1), 1–162. <https://doi.org/10.1186/s40359-021-00669-5>
- Abu Baker, J. (2021, June 10). Up to 5 in a group allowed from Jun 14; dining-in may resume on Jun 21 in phased easing of COVID-19 curbs. *Channel News Asia*. <https://www.channelnewsasia.com/singapore/cap-of-5-people-social-gatherings-dining-in-phase-3-covid-19-1846791>
- Adler, N. E., Epel, E. S., Castellazzo, G., & Ickovics, J. R. (2000). Relationship of subjective and objective social status with psychological and physiological functioning: Preliminary data in healthy white women. *Health Psychology*, 19(6), 586–592. <https://doi.org/10.1037/0278-6133.19.6.586>
- Allahyari, T., Saraji, G. N., Adi, J., Hosseini, M., Iravani, M., Younesian, M., & Kass, S. J. (2019). Cognitive failures, driving errors, and driving accidents. *International Journal of Occupational Safety and Ergonomics*, 14(2), 149–158. <https://doi.org/10.1080/10803548.2008.11076759>
- Almeida, D. M., Wethington, E., & Kessler, R. C. (2002). The daily inventory of stressful events: An interview-based approach for measuring daily stressors. *Assessment (Odessa, Fla.)*, 9(1), 41–55. <https://doi.org/10.1177/1073191102009001006>
- Ang, H. M. (2021, July 20). Return to Phase 2 (Heightened Alert): Dining-in to be suspended, group sizes back down to 2. *Channel News Asia*. <https://www.channelnewsasia.com/singapore/phase-2-heightened-alert-dining-in-gathering-group-size-covid-19-2047211>
- Arıcan, H., & Kafadar, H. (2022). Examining the relationship between coronavirus anxiety and cognitive failures. *Neuropsychiatric Investigation*, 60(4), 1–7. <https://doi.org/10.5152/NeuropsychiatricInvest.2022.22012>
- Asmundson, G. S. G., Rachor, G., Drakes, D. H., Boehme, B. A. E., Paluszek, M. M., & Taylor, S. (2022). How does COVID stress vary across the anxiety-related disorders? Assessing factorial invariance and changes in COVID Stress Scale scores during

- the pandemic. *Journal of Anxiety Disorders*, 87, 102554. <https://doi.org/10.1016/j.janxdis.2022.102554>
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1). <https://doi.org/10.18637/jss.v067.i01>
- Ben-Shachar, M., Lüdtke, D., & Makowski, D. (2020). Effect size: Estimation of effect size indices and standardized parameters. *Journal of Open Source Software*, 5(56), 2815. <https://doi.org/10.21105/joss.02815>
- Bolger, N., Stadler, G., & Laurenceau, J.-P. (2012). Power analysis for intensive longitudinal studies. In *Handbook of research methods for studying daily life* (pp. 285–301). The Guilford Press.
- Bridger, R. S., Johsen, SÅK, & Brasher, K. (2013). Psychometric properties of the Cognitive Failures Questionnaire. *Ergonomics*, 56(10), 1515–11524. <https://doi.org/10.1080/00140139.2013.821172>
- Broadbent, D. E., Cooper, P. F., FitzGerald, P., & Parkes, K. R. (1982). The Cognitive Failures Questionnaire (CFQ) and its correlates. *British Journal of Clinical Psychology*, 21(1), 1–16. <https://doi.org/10.1111/j.2044-8260.1982.tb01421.x>
- Burdett, B. R. D., Charlton, S. G., & Starkey, N. J. (2016). Not all minds wander equally: The influence of traits, states and road environment factors on self-reported mind wandering during everyday driving. *Accident Analysis & Prevention*, 95(Pt A), 1–7. <https://doi.org/10.1016/j.aap.2016.06.012>
- Carlander, A., Lekander, M., Asmundson, G. J. G., Taylor, S., Bagge, R. O., & Bagge, A.-S. L. (2022). COVID-19 related distress in the Swedish population: Validation of the Swedish version of the COVID Stress Scales (CSS). *PloS One*, 17(2), e0263888. <https://doi.org/10.1371/journal.pone.0263888>
- Chia, J. L., & Hartanto, A. (2021). Cognitive barriers to COVID-19 vaccine uptake among older adults. *Frontiers in Medicine*, 8, 756275. <https://doi.org/10.3389/fmed.2021.756275>
- Colom, R., Martínez-Molina, A., Shih, P. C., & Santacreu, J. (2010). Intelligence, working memory, and multitasking performance. *Intelligence*, 38(6), 543–551. <https://doi.org/10.1016/j.intell.2010.08.002>
- Goh, A. Y. H., Chia, S. M., Majeed, N. M., Chen, N. R. Y., & Hartanto, A. (2023). Untangling the additive and multiplicative relations between natural scenery exposure and human–animal interaction on affective well-being: Evidence from daily diary studies. *Sustainability*, 15(4), 2910. <https://doi.org/10.3390/su15042910>
- Hartanto, A., & Yang, H. (2020). Testing theoretical assumptions underlying the relation between anxiety, mind wandering, and task-switching: A diffusion model analysis. *Emotion*, 22(3), 493–510. <https://doi.org/10.1037/emo0000935>
- Jorgensen, T. D., Pornprasertmanit, S., Schoemann, A. M., & Rosseel, Y. (2021). *semTools: Useful tools for structural equation modeling* (0.5-5) [R]. <https://CRAN.R-project.org/package=semTools>
- Kane, M. J., Hambrick, D. Z., & Conway, A. R. A. (2005). Working memory capacity and fluid intelligence are strongly related constructs: Comment on Ackerman, Beier, and Boyle (2005). *Psychological Bulletin*, 131(1), 66–71. <https://doi.org/10.1037/0033-2909.131.1.66>
- Khosravani, V., Asmundson, G. J. G., Taylor, S., Sharifi Bastan, F., & Samimi Ardestani, S. M. (2021). The Persian COVID Stress Scales (Persian-CSS) and COVID-19-related stress reactions in patients with obsessive-compulsive and anxiety disorders. *Journal of Obsessive-Compulsive and Related Disorders*, 28, 100615–100615. <https://doi.org/10.1016/j.jocrd.2020.100615>
- Klein, K., & Boals, A. (2001). The relationship of life event stress and working memory capacity. *Applied Cognitive Psychology*, 15(5), 565–579. <https://doi.org/10.1002/acp.727>
- Knowles, J. E., Frederick, C., & Whitworth, A. (2020). *merTools: Tools for Analyzing Mixed Effect Regression Models* (Version 0.5.2). <https://cran.r-project.org/package=merTools>
- Kottwitz, M. U., Wehrt, W., Gerhardt, C., Augusto Coelho, D., Schmutz, D., & Elfering, A. (2022). Yesterday’s work–home conflict and actigraphically recorded sleep-onset latency as predictors of today’s cognitive failure. *Journal of Business and Psychology*, 37(3), 509–524. <https://doi.org/10.1007/s10869-021-09766-z>
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in linear mixed effects models. *Journal of Statistical Software*, 82(13). <https://doi.org/10.18637/jss.v082.i13>
- Lange, S., & Süß, H. M. (2014). Measuring slips and lapses when they occur—Ambulatory assessment in application to cognitive failures. *Consciousness and Cognition*, 24, 1–11. <https://doi.org/10.1016/j.concog.2013.12.008>
- Larson, G. E., & Merritt, C. R. (1991). Can accidents be predicted? An empirical test of the Cognitive Failures Questionnaire. *Applied Psychology: An International Review*, 40(1), 37–45. <https://doi.org/10.1111/j.1464-0597.1991.tb01356.x>
- Maas, C. J. M., & Hox, J. J. (2004). The influence of violations of assumptions on multilevel parameter estimates and their standard errors. *Computational Statistics & Data Analysis*, 46(3), 427–440. <https://doi.org/10.1016/j.csda.2003.08.006>
- Mahoney, A. M., Dalby, J. T., & King, M. C. (1998). Cognitive failures and stress. *Psychological Reports*, 82(3_suppl), 1432–1434. <https://doi.org/10.2466/pr0.1998.82.3c.1432>
- Majeed, N. M., Kasturiratna, K. T. A. S., Lua, V. Y. Q., Li, M. Y., & Hartanto, A. (2023). Discrimination and cognitive failures in Singapore and the US: An investigation of between- and within-persons associations through multilevel modelling. *International Journal of Intercultural Relations*, 94, 101805. <https://doi.org/10.1016/j.ijintrel.2023.101805>
- Majeed, N. M., Tan, J. J. X., Tov, W., & Hartanto, A. (2021). Dispositional optimism as a buffer against emotional reactivity to daily stressors: A daily diary approach. *Journal of Research in Personality*, 93, 104105. <https://doi.org/10.1016/j.jrp.2021.104105>
- Mathieu, E., Ritchie, H., Rodés-Guirao, L., Appel, C., Gavrilow, D., Giattino, C., Hasell, J., Macdonald, B., Dattani, S., Beltekian, D., Ortiz-Ospina, E., & Roser, M. (2020). *Coronavirus Pandemic (COVID-19)*. Our World In Data. <http://OurWorldInData.org>
- McVay, J. C., & Kane, M. J. (2010). Does mind wandering reflect executive function or executive failure? Comment on Smallwood and Schooler (2006) and Watkins (2008). *Psychological Bulletin*, 136(2), 188–197. <https://doi.org/10.1037/a0018298>
- Morin, A., & Racy, F. (2021). Dynamic self-processes. In J. F. Rauthmann (Ed.), *The handbook of personality dynamics and processes* (pp. 365–386). <https://doi.org/10.1016/B978-0-12-813995-0.00015-7>
- Muthén, B. O., & Muthén, L. K. (2012). *Mplus version 7: User’s guide*. Author.
- Ng, M. H. S., Lua, V. Y. Q., Majeed, N. M., & Hartanto, A. (2022). Does trait self-esteem serve as a protective factor in

- maintaining daily affective well-being? Multilevel analyses of daily diary studies in the US and Singapore. *Personality and Individual Differences*, *198*, 111804. <https://doi.org/10.1016/j.paid.2022.111804>
- Ng, T. L. Y., Majeed, N. M., Lua, V. Y. Q., & Hartanto, A. (2023). Do executive functions buffer against COVID-19 stress? A latent variable approach. *Current Psychology*. <https://doi.org/10.1007/s12144-023-04652-8>
- Noël, M.-P. (2009). Counting on working memory when learning to count and to add: A preschool study. *Developmental Psychology*, *45*(6), 1630–1643. <https://doi.org/10.1037/a0016224>
- Pakpour, A. H., Griffiths, M. D., & Lin, C.-Y. (2021). Assessing psychological response to the COVID-19: The fear of COVID-19 Scale and the COVID Stress Scales. *International Journal of Mental Health and Addiction*, *19*(6), 2407–2410. <https://doi.org/10.1007/s11469-020-00334-9>
- Pulido, E. G., & Jimenez, L. (2020). *Validation to Spanish version of the COVID-19 Stress Scale* [Preprint]. PsyArXiv. <https://doi.org/10.31234/osf.io/qgc3h>
- R Core Team. (2020). R: A language and environment for statistical computing (3.6.3) [Computer software]. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Revelle, W. (2021). *psych: Procedures for psychological, psychometric, and personality research* (2.1.6) [R]. Northwestern University.
- Rush, J., Rast, P., Almeida, D. M., & Hofer, S. M. (2019). Modeling long-term changes in daily within-person associations: An application of multilevel SEM. *Psychology and Aging*, *34*(2), 163–176. <https://doi.org/10.1037/pag0000331>
- Smallwood, J., & Schooler, J. W. (2006). The restless mind. *Psychological Bulletin*, *132*(6), 946–958. <https://doi.org/10.1037/0033-2909.132.6.946>
- Smeekens, B. A., & Kane, M. J. (2016). Working memory capacity, mind wandering, and creative cognition: An individual-differences investigation into the benefits of controlled versus spontaneous thought. *Psychology of Aesthetics, Creativity, and the Arts*, *10*(4), 389–415. <https://doi.org/10.1037/aca0000046>
- Taylor, S., Landry, C. A., Paluszek, M. M., Fergus, T. A., McKay, D., & Asmundson, G. J. G. (2020). Development and initial validation of the COVID Stress Scales. *Journal of Anxiety Disorders*, *72*, 102232. <https://doi.org/10.1016/j.janxdis.2020.102232>
- Unsworth, N., Brewer, G. A., & Spillers, G. J. (2012a). Variation in cognitive failures: An individual differences investigation of everyday attention and memory failures. *Journal of Memory and Language*, *67*(1), 1–16. <https://doi.org/10.1016/j.jml.2011.12.005>
- Unsworth, N., McMillan, B. D., Brewer, G. A., & Spillers, G. J. (2012b). Everyday attention failures: An individual differences investigation. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *38*(6), 1765–1772. <https://doi.org/10.1037/a0028075>
- Vatansever, D., Wang, S., & Sahakian, B. J. (2021). COVID-19 and promising solutions to combat symptoms of stress, anxiety and depression. *Neuropsychopharmacology*, *46*(1), 217–218. <https://doi.org/10.1038/s41386-020-00791-9>
- Wallace, J. C., & Vodanovich, S. J. (2003a). Can accidents and industrial mishaps be predicted? Further investigation into the relationship between cognitive failure and reports of accidents. *Journal of Business and Psychology*, *17*(4), 503–514. <https://doi.org/10.1023/A:1023452218225>
- Wallace, J. C., & Vodanovich, S. J. (2003b). Workplace safety performance: Conscientiousness, cognitive failure, and their interaction. *Journal of Occupational Health Psychology*, *8*(4), 316–327. <https://doi.org/10.1037/1076-8998.8.4.316>
- Wickham, H., François, R., Henry, L., Müller, K., & Vaughan, D. (2022). *dplyr: A Grammar of data manipulation* (1.1.0) [R]. <https://CRAN.R-project.org/package=dplyr>
- World Health Organization. (2022). *WHO Coronavirus (COVID-19) Dashboard*. World Health Organization. <https://covid19.who.int>
- Xu, C., Xu, Y., Xu, S., Zhang, Q., Liu, X., Shao, Y., Xu, X., Peng, L., & Li, M. (2020). Cognitive reappraisal and the association between perceived stress and anxiety symptoms in COVID-19 isolated people. *Frontiers in Psychiatry*, *11*, 858. <https://doi.org/10.3389/fpsy.2020.00858>
- Yu, Q., & Li, B. (2022) *mlma: Multilevel mediation analysis* (6.2-1) [R]. <https://CRAN.R-project.org/package=mlma>